



Top Surf[®]

Nonionic 80/20 Surfactant

ACTIVE INGREDIENTS

Alkylpolyethoxyalkylene ethers and other ethoxylated derivatives	80.0%
--	-------

INERT INGREDIENTS	<u>20.0%</u>
-------------------------	--------------

TOTAL	100.0%
-------------	--------

STOP-READ LABEL BEFORE USING.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Do not store near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

Causes eye irritation. Avoid contact with eyes. In case of contact, flush eyes with water for at least 15 minutes and get medical attention.

If taken internally, call a physician at once.

STORAGE AND DISPOSAL

Do not use, pour, spill, or store near heat or open flame. Use adequate ventilation when handling.

Do not reuse empty container. Triple rinse, or equivalent, then offer for recycling or for disposal in a sanitary landfill, or by other means approved by State and local authorities.

NET CONTENTS

Distributed By:
Winfield Solutions, LLC
P.O. Box 64589, St. Paul, MN 55164-0589

0/0511/9

DIRECTIONS FOR USE

TOP SURF is a water soluble surfactant designed for use in certain agricultural and horticultural sprays.

Suggested rates for use with various categories of chemicals are as follows:

Chemical Group	Pints of TOP SURF to Add Per 100 Gallons of Spray Mixture
Insecticides, Miticides & Fungicides	0.5 to 2 pints
Herbicides	1 to 4 pints
Glyphosate Formulations Without Added Surfactants	8 pints
Defoliant & Desiccants	1 to 4 pints
Fertilizers & Micronutrients	0.5 to 2 pints
Livestock Sprays	0.25 to 0.5 pints

Note: Carefully read and follow the directions contained on the label of the herbicide, insecticide, or livestock spray with which TOP SURF is to be used. For local recommendations consult a competent agricultural authority.

Cotton Pickers: Use about 4 to 6 ounces per 50 gallons of water, or 2 to 3 ounces per 25 gallon picker tank. While filling tank with water, add TOP SURF when about one-half full. Mix thoroughly with water by circulation or by stirring.

Gins: Use at the rate of 6 ounces of TOP SURF to 50 gallons of water for restoring moisture to cotton.

Aerial Application: When applying aerially, follow pesticide label directions including minimum water volume per acre. Do not substitute TOP SURF for water required by the pesticide label.

Aquatic Application: Apply at the rate recommended as described by pesticide label.

Notice of Warranty

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, NOR IS ANY REPRESENTATIVE OF SELLER AUTHORIZED TO MAKE ANY SUCH WARRANTY OR MODIFY THESE TERMS. This warranty does not extend to the storage, handling or use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such storage, handling or use. Seller shall not be responsible for incidental or consequential damages, if any, resulting from a breach of warranty.

TOP SURF® is a registered trademark of Winfield Solutions, LLC.